

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
http://ageconsearch.umn.edu
aesearch@umn.edu

Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

Analysis on Current Situations of Endangered *P. intermedius* S. Y. Chen

XU Yuan-ke¹*, ZHANG Shao-hua¹, PENG Xue-ku², LIU Rao³

1. Jingning She Autonomous County Forestry Bureau, Jingning 323500, China; 2. Jingnan Township Forestry Workstation of Jingning She Autonomous County, Jingning 323500, China; 3. Science and Technology Service Department of Jingning She Autonomous County Bureau of Science and Technology, Jingning 323500, China

Abstract We carried out a general survey of current situations of *P. intermedius* S. Y. Chen. Results show that *P. intermedius* S. Y. Chen is distributed in Dongkeng Township, Jingnan Township and Yanxi Township. It covers an area of 262.51 hm². The distribution areas are narrow and limited. What's worse, it suffers from serious destruction and has been withered in large areas. Besides, according to current introduction test, it is difficult to survive in other places. In accordance with *Exploring the Criteria for Endangered Categories of Bamboo Species from China* put forward by Ma Naixun *et al* from the Research Institute of Subtropical Forestry of Chinese Academy of Forestry (RISFCAF), we propose incorporating *P. intermedius* S. Y. Chen into vulnerable species for protection.

Key words P. intermedius S. Y. Chen, Current situations, Analysis on endangered situations

P. intermedius S. Y. Chen, also called Lishi Bamboo, has tasty bamboo shoots. It is as high as 3 to 4 m with diameter 1 to 2 cm. New bamboo takes on green covered with white powder. At upper part of bamboo joint, there are small rough hairs which are not easily visible but vulnerable to fall off. Sheaths of bamboo shoots fall late and the length between joints is nearly equal. After rough hairs falling off, upper edge will become shriveled. It has slightly grown sheaths and lacks auricles. Sheath ligule is in concave or arc shape with height of 3 to 4 mm. Leaves of sheaths, like needle, open or overturn, and slightly roll towards backside. Leaves are generally 12 to 25 cm long and 25 to 32 cm wide. The bamboo-shoot sprouts from the late ten days of April. Bamboo poles are stiff and suitable for umbrella handle, but hard to split[1]. As a rare species, P. intermedius S. Y. Chen is unique in Jingning She Autonomous County. Bamboo-shoot sprouts from the late ten days of April to the first and middle ten days of May. This bamboo shoot is fragrant and sweet, its lower part is more fresh and tender then upper part. Small bamboo shoots pickled in Dongkeng area of Jingning She Autonomous County are well received by town dwellers and citizens. Main materials of these pickled shoots are P. intermedius S. Y. Chen. Unfortunately. large area of bamboo grove is found withered in 1997^[2]. To enhance protection of P. intermedius S. Y. Chen, we carried out survey for this resource and here analyze its endangered situation.

1 Survey method

Through visiting, we got preliminary understanding of the distribution range of *P. intermedius* S. Y. Chen. According to

similarity of environment site, we expanded the survey areas. After knowing its distribution range, we surveyed its area, growth condition and elevation range of natural distribution area using sampling survey and topographic map plotting method.

2 Survey results

The *P. intermedius* S. Y. Chen is mainly distributed in Dongkeng Township, Jingnan Township and Yanxi Township of Jingning County, with an area of 262.51 hm² (shown in Table 1).

3 Criteria for judging endangering and analysis

3.1 Criteria for judging endangering According to Exploring the Criteria for Endangered Categories of Bamboo Species from China put forward by Ma Naixun et al from the Research Institute of Subtropical Forestry of Chinese Academy of Forestry, we selected the ecological amplitude, distribution range, population quantity, value of development and utilization, and endemic species to divide the protection of bamboo species in China into endangered, rare and vulnerable levels. Then, we analyzed endangered situations of *P. intermedius* S. Y. Chen. Endangered bamboo species. This type of bamboo species is distributed in a narrow area with small ecological amplitude, the genetic feature is highly special and storage in strange place is hard to succeed under general technical conditions. Alternatively, among lots of types of bamboo species, this has important or unique economic, development and utilization value. Nevertheless, in recent two or three decades, resource damage is serious, its growth condition becomes extremely bad and the situation is worsening. Another case is autogenus that has important value or potential value in science and needs special protection[3].

Received: January 29, 2011 Accepted: February 20, 2012

^{*} Corresponding author. E-mail: jnxyk026@126.com

3.1.2 Rare species. The distribution range is narrow, by now, *P. intermedius* S. Y. Chen is found only in one county and district. In adjacent counties or districts, only little population has been found. Generally, the population area is less

than 20 hm² with serious artificial disturbance. Alternatively, it has certain economic and scientific value and should receive close attention^[3].

Table 1 Distribution of P. intermedius S. Y. Chen

Township	Village	Distribution	Elevation	Area	Main associated plants
		points	m	hm²	
Dongkeng	Makeng	4	640 -830	1.73	Pine, China fir, oak, Schima superba, and azalea
	Zhubu	3	700 -850	2.67	Pine, oak, Qinggang oak, etc.
	Hecun	2	600 -840	7.00	Pine, oak, Schima superba, etc.
	Beishan	1	500 -520	0.20	Oak, Castanopsis sclerophylla, azalea, etc.
	Baihe	3	530 -800	6.40	Pine, oak, P. tomentosa, azalea, etc
	Taoyuan	5	550 - 760	11.00	Pine, oak, Schima superba, Alnus cremastogyne, etc.
	Zhangkeng	2	500 -650	8.00	Pine, China fir, bamboo, azalea, etc.
	Tangbei	1	610 - 690	2.00	Pine, China fir, oak, azalea, Alnus cremastogyne, etc
	Dazhangkeng	3	750 – 1 000	3.33	Pine, China fir, oak, azalea, Castanopsis sclerophylla, etc
	Wushantou	1	730 – 780	2.00	Pine, China fir, oak, azalea, etc
	Gendixian	1	700 -800	3.33	Pine, oak, Castanopsis sclerophylla, azalea, etc.
	Xulong	1	620 - 750	4.67	Pine, oak, Castanopsis sclerophylla, azalea, etc.
	Dongkeng	1	600 - 760	1.33	Pine, China fir, Castanopsis sclerophylla, azalea, etc.
Jingnan	Dongtang	11	870 – 980	19.00	Pine, China fir, Dicranopteris linearis, Schima superba, and azalea
	Zhongxi	5	900 - 940	7.33	Pine, China fir, Castanopsis sclerophylla, and azalea
	Chengtian	11	650 -850	16.53	Pine, China fir, Castanopsis sclerophylla, and azalea
	Miaokeng	6	700 –750	20.33	Pine, China fir, Castanopsis sclerophylla, azalea, and small bamboo
Yanxi	Huangtong	10	700 –750	35.00	Pine, China fir, Castanopsis sclerophylla, and azalea
	Shiti	15	600 - 750	36.33	Pine, China fir, Castanopsis sclerophylla, azalea, and small bamboo
	Yanxi	14	520 -600	41.33	Pine, China fir, Castanopsis sclerophylla, and Schima superba
	Dongshan	3	650 -800	4.67	Pine, China fir, Castanopsis sclerophylla, Schima superba, and
					Dicranopteris linearis
	Meiniao	8	600 - 750	21.33	Pine, China fir, Castanopsis sclerophylla, Schima superba, and
					Dicranopteris linearis
	Zhewan	2	600 -650	3.33	Pine, China fir, Castanopsis sclerophylla, and oak
	Daqiutian	4	530 -700	3.67	Pine, China fir, oak, Alnus cremastogyne and P. tomentosa

- **3. 1. 3** Vulnerable bamboo species. The distribution of this species is similar to rare species. Although there is large distribution in many districts and counties, the area is little and total area is less than 100 hm². This species has certain development and utilization value, but along with excessive development and ecological deterioration, it probably becomes the endangered species^[3].
- **3.2** Analysis on endangered situation The *P. intermedius* S. Y. Chen is mainly distributed in Dongkeng Township, Jingnan Township and Yanxi Township of Jingning County, with an area of 262.51 hm². Although it is larger than 100 hm², its distribution range is very narrow and large area of *P. intermedius* S. Y. Chen has once been withered. What's worse, in recent years, some of *P. intermedius* S. Y. Chen are found flower-

ing, and artificial destruction is serious. Besides, according to current introduction test, it is difficult to survive in other places. Therefore, we propose incorporating *P. intermedius* S. Y. Chen into vulnerable species for protection.

References

- [1] ZHU SL, MA NX, FU MY.) Flora of Chinese bamboo[M]. Beijing: China Forestry Publishing House, 1994. (in Chinese).
- [2] SHEN X, WANG KH, ZHOU XL, et al. New century bamboo development research of Zhejiang [M]. Beijing: China Agricultural Science and Technology Press, 2001; 257 –258. (in Chinese).
- [3] MA NX, CHEN GC, ZHANG WY. Exploring the criteria for endangered categories of bamboo species from China[J]. Scientia Silvae Sinicae, 2006, 42(6): 56 -60. (in Chinese).