



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



Adoption of improved quinoa varieties among smallholder farmers in the Peruvian Andes

Cindybell GAMBOA, Goedele VAN DEN BROECK and Miet MAERTENS

Bioeconomics Working Paper Series

Working Paper 2018/7



KU LEUVEN

Division of Bioeconomics
Department of Earth and Environmental Sciences
University of Leuven
Geo-Institute
Celestijnenlaan 200 E – box 2411
3001 Leuven (Heverlee)
Belgium
<http://ees.kuleuven.be/bioecon/>

Adoption of improved quinoa varieties among smallholder farmers in the Peruvian Andes

Cindybell GAMBOA ^{1, 2}, Goedele VAN DEN BROECK ¹ and Miet MAERTENS ¹

Abstract

Quinoa production has boomed tremendously in the Andes since the early 2010s. It is widely promoted as a highly nutritious food and is rapidly changing from a subsistence crop into a high-value cash crop. This entails opportunities for upward income mobility and improved food security for Andean smallholders. However, farmers in the Andean highlands cannot reap the full benefits of lucrative market opportunities due to poor agricultural technologies and low quinoa productivity levels. While investments into developing improved quinoa varieties are ongoing, breeding programs are not tailored to the needs of smallholder quinoa farmers producing for own consumption, the local market and the export market. In this paper, we study farmers' preferences and willingness to pay for improved quinoa varieties in the Junín region in Peru. We use data from a choice experiment among 458 Andean smallholders and estimate generalized multinomial logit models to control for preference and scale heterogeneity. We find that farmers generally prefer improved varieties over traditional varieties, with mildew-resistance as most important crop trait. This is in line with mildew being a serious constraint to quinoa production in the Andes. In general, farmers prefer varieties that are characterized by larger grain sizes, higher yield levels, lower levels of saponin, and a reduced maturation period. Yet, food-insecure farmers are found to be indifferent to early maturity and a larger grain size, which can be explained by a lower degree of commercialization among these farmers. Our results imply that developing mildew-tolerant and higher-yielding varieties with a medium to low saponin content is a priority if investments in quinoa technologies is to benefit small-scale and food insecure farmers in the Andean highlands of Peru.

Key Words: Quinoa, choice experiment, scale and preference heterogeneity, generalized multinomial logit, Peru

JEL classification: O13, O33, Q12, Q16

Corresponding author: Cindybell.gamboa@kuleuven.be

Acknowledgements

We are thankful to José Falconi for useful comments on the identification of the attributes and attribute levels that are relevant for the adoption of improved quinoa varieties and to the Dirección Regional de Agricultura and the Agencias Agrarias for supporting the data collection in Junín region in Peru.

¹ Division of Bioeconomics, Department of Earth and Environmental Sciences, KU Leuven, Celestijnenlaan 200E – box 2411, 3001 Leuven, Belgium.

² Universidad Nacional Agraria La Molina, Av. La Molina s/n, La Molina, Perú

Adoption of improved quinoa varieties among smallholder farmers in the Peruvian Andes

1. Introduction

Quinoa is widely promoted as a highly nutritious food and demand from high-income countries is rapidly increasing (Lester, 2006; Ruiz et al., 2014). As a consequence, production of quinoa has boomed tremendously in the Andes region since 2010, with Peru as main producing and exporting country. While quinoa is traditionally cultivated as a common staple and subsistence crop by smallholders in the Andes region, it is increasingly produced by large-scale farms in the Peruvian coastal areas as a high-value export crop (Ofstehage, 2012). In addition, the number of countries growing quinoa outside the Andean region, like the United States of America, United Kingdom and Denmark among others, has risen sharply since 2014 (Bazile, Pulvento, et al., 2016; Bazile, Bertero, & Nieto, 2015; Zurita-Silva, Fuentes, Zamora, Jacobsen, & Schwember, 2014). This transition has been supported by recent investments into the development of improved quinoa varieties, focusing on higher yields, a reduced maturation period and a sweeter taste (Zurita-Silva et al., 2014).

It is unlikely that the current developments towards improved quinoa varieties will benefit small-scale producers and improve local food security in Andean regions. Investments are mainly targeted to the export sector and large-scale producers in coastal areas. Smallholders face different constraints to crop production and marketing than large-scale producers and might have different preferences for certain traits of improved quinoa varieties. For example, taste preference in Andean regions where quinoa belongs to the traditional diet may differ from taste preferences in international markets where quinoa has become a healthy food hype. In addition, one of the major impediments to reaching high productivity in the Andes region is quinoa's susceptibility to mildew, a plant disease caused by fungi (Sven-Erik Jacobsen, 2003; Ruiz et al., 2014). This pathogen causes the development of chlorotic and necrotic spots in the leaves and premature leaf fall, leading to yield reductions of up to 99% in susceptible cultivars (Danielsen & Munk, 2004; Zurita-Silva et al., 2014). Given that smallholders in the Andes region still cultivate almost 90% of the total area allocated to quinoa production in Peru, the development of mildew-resistant varieties is thus a key priority for researchers and policy makers.

In order to increase adoption rates of improved varieties among quinoa producers, it is crucial to take into account their specific requirements and needs ex-ante, before new varieties are developed and introduced (Sibiya, Tongoona, Derera, & Makandaa, 2013). In this paper,

we study farmers' preferences and willingness to pay for improved quinoa varieties in the Junín region in Peru, using a choice experiment among 458 Andean smallholders. A choice experiment is a stated preference method to assess which specific traits of a certain hypothetical scenario (in this case improved quinoa varieties) are valued most. Previous research has used choice experiments to investigate the adoption of improved seed varieties of different staple food crops, such as maize in Zimbabwe (Kassie et al., 2017) and Mexico (Sánchez-Toledano, Kallas, & Gil-Roig, 2017), pearl millet in India (Smale, Diressie, & Birol, 2016), beans in Burundi (Lambrecht, Vranken, Merckx, Vanlauwe, & Maertens, 2015), rice in India (Ward, Ortega, Spielman, & Singh, 2014), and sorghum and teff in Ethiopia (Asrat, Yesuf, Carlsson, & Wale, 2010). These studies show that farmers' preferences for improved staple crop seeds are heterogeneous and depend on the local context, and demonstrate the usefulness of using choice experimental research to provide ex-ante insights on farmers' preferences for agricultural technologies. To the best of our knowledge, no studies have analyzed the preferences of Andean quinoa farmers for improved quinoa seeds. This paper thereby makes a relevant contribution as quinoa is an increasingly important crop in local as well as international markets. For smallholder Andean quinoa farmers to benefit fully from lucrative market opportunities and ongoing investments in quinoa research and technology development, it is imperative to understand their preferences. In addition, we use an approach that allows us to analyze preference heterogeneity as well as scale heterogeneity while most existing choice experiment studies do not distinguish between these two sources of heterogeneity.

The paper is structured as follows. The second section provides background information on quinoa production and the development of improved varieties in Peru. The third section describes the data and methodology, while the fourth section presents the results. In the fifth section we discuss the results and the sixth section concludes.

2. Background

2.1 Export and production of quinoa

The international demand for quinoa as a highly nutritious food has increased rapidly since 2007 (MINAGRI, 2017). Peru has consolidated its position as market leader in quinoa exports, bypassing Bolivia to be the largest quinoa exporter since 2014. The export volume in Peru increased with an average annual growth rate of 37% during the period 2000-2017 (Figure 1). At the same time, the export price in Peru increased considerably, from 1.27 USD/kg on average in 2000 to 2.34 USD/kg in 2017, corresponding to an average annual price increase of 3.69%.

The sharpest increase is observed between 2008 and 2014 when the price jumped from 2.42 to 5.42 USD/kg. After 2014, the price decreased again to a similar level as in 2008.




Figure 1. Export volume and export price of Peruvian quinoa

Source: Authors’ elaboration from Ministry of Agriculture and Irrigation (MINAGRI), over 2000-2017.

To meet this international demand, the area allocated to quinoa production has increased tremendously in Peru. Quinoa is cultivated in 18 out of the 24 regions in Peru (Apaza, Cáceres, Estrada, & Pinedo, 2015). An important distinction can be made between the six traditional producer-regions in the Andes (Apurimac, Ayacucho, Cajamarca, Cusco, Junín and Puno), which account for 89% of national quinoa area in 2016, and 12 secondary and new producer-regions (Arequipa, Ancash, Huancavelica, Huánuco, Ica, Lambayeque, La Libertad, Lima, Moquegua, Pasco, Piura and Tacna), which account for 11% of the national area (Figure 2). These new producer-regions are mainly located in the coastal agro-industrial zones of Peru, where the government has actively promoted the expansion of quinoa cultivation since 2014. Because of the large adaptability of quinoa to other environmental conditions and a more efficient water and soil use, productivity in these areas is higher than in the traditional producer-regions; with average yields of 1.9 ton/ha, compared to 1.3 ton/ha in traditional regions (Bedoya-Perales, Pumi, Mujica, Talamini, & Padula, 2018).




Figure 2. Evolution of quinoa area harvested and production in Peru in traditional quinoa-producing (TP) regions and secondary and new producing (SNP) regions.

Source: Authors' elaboration from MINAGRI, over 1991-2016

In the traditional producer-regions in the Andes, quinoa production is constrained by adverse climate and agro-ecological conditions such as drought, frost, wind, hail, soil salinity and poor soil fertility (S.-E. Jacobsen, Mujica, & Jensen, 2003; S. E. Jacobsen, Mujica, & Ortiz, 2003; Sven-Erik Jacobsen, 2003). Also the occurrence of mildew, the most important plant disease in quinoa production, can seriously constraint productivity (Sven-Erik Jacobsen, 2003; Ruiz et al., 2014). The maturation period for quinoa is relatively long – six months for the most common variety used in the research area – which is a limiting factor for farmers seeking several crop rotations per year. In addition, farmers face constraints to meet requirements in export markets due to a high saponin content in the seed hull, and small and non-uniform grain sizes (Bazile, Jacobsen, & Verniau, 2016; Bedoya-Perales et al., 2018).

2.2 Development of improved varieties

The current development of improved quinoa varieties mostly focuses on production for the international market. In Europe and North America, breeding programs concentrate on early maturity, high yield, uniformity of seeds and sweet cultivars for commercial production (Bazile, Jacobsen, et al., 2016). In Peru, the Cereal and Native Grain Research Program at the National Agricultural University La Molina is developing genetically improved quinoa seeds, with a focus on early maturation and reduction in plant height for easier introduction of quinoa in modern agricultural systems (Gomez-Pando & Eguiluz-de la Barra, 2013).

Recently, there has been an effort to develop improved varieties for the traditional producer-regions in Peru. Nationally, 35 quinoa varieties have been identified, of which eight are improved (Bojanic, 2011; Mujica, 1992). The improved varieties have been obtained

through conventional breeding methods, such as selection and hybridization, by the experimental stations of the National Institute of Agricultural Research (INIA) (Apaza et al., 2015). Breeding programs have mainly focused on tolerance to plagues and diseases, high yield, and large grain size. The varieties have been developed for specific producer-regions, such as Puno (INIA 420-Negra Collana, INIA 415-Pasankalla, Salcedo INIA, Ilpa INIA and INIA 431-Altiplano), Cusco (Quillahuaman INIA and INIA 427-Amarilla Sacaca) and Junín (INIA 433-SantaAna/AIQ/FAO). The uptake of these improved varieties among smallholder farmers is rather limited. This could be related to a lack of farmer awareness about improved quinoa varieties; a high cost of improved varieties combined with capital constraints and risk aversion among farmers; or the traits of the improved varieties not being in line with farmers' preferences.

2.3 Quinoa and food security

Quinoa production and trade entail important opportunities for upward income mobility and improved food security among Andean smallholders (S.-E. Jacobsen et al., 2003). Quinoa is one of the main grain crops supplying highly nutritious food for farm-households in the Andean region and also has the potential to contribute to food security in other regions of the developing world (Sven-Erik Jacobsen, 2003). Being a strategic Andean crop in the region, quinoa has attracted political and research interest since 1999. The Peruvian government has been promoting quinoa cultivation to improve the food security of the impoverished Andean population through various social programs (Bedoya-Perales et al., 2018; Hellin & Higman, 2005). In 2008, the food assistance program “National Food Support Program - PRONAA” was authorized by the state to purchase Andean agricultural products, with quinoa being one of those products, directly from local small farmers. In 2012, the National Program School Food Qali Warma provides a quality food service, appropriate to local consumption habits to girls and boys who are enrolled in public educational institutions at the primary and kindergarten levels, in which the program incorporates quinoa as a processed product.

The introduction and adoption of improved quinoa varieties could have a positive impact on improving food security and reducing poverty in the Andean region. Farmers might benefit from a higher productivity while consumers might favor a better taste. However, a study for Puno showed that a change in quinoa's taste did not lead to a substantial difference in households' nutrition, because only a relatively small share of households' overall diets consists of quinoa (Stevens, 2017).

3. Data and methodology

3.1 Research area

In this paper, we focus on traditional quinoa producers in the Junín region, located in the central highlands of Peru. In 2012, this region was considered to be moderately vulnerable to food insecurity (MIDIS, 2012). Before 2013, quinoa was a subsistence crop for smallholder farmers and was mainly produced on small plots of land for self-consumption (INEI, 2013) (Table A1). Andean farmers usually diversify crop production with potato, barley and maize as the most common crops in the Junín region (DRAJ, 2017). Quinoa is cultivated in two agro-ecological zones in the Andean region of Junín, namely Quechua and Suni zones with altitudes between 2300-3500 m.a.s.l. and 3400-3800 m.a.s.l., respectively, and average annual temperature around 11-16°C and annual rainfall between 500 and 1200 mm (Tapia & Fries, 2007).

The quinoa production in this region is highly prone to mildew (*Peronospora farinosa*) (Rojas et al., 2015). It affects the germination and the formation of the first sheet of quinoa and causes a risk of yield loss. The Junín region is one of the five sub-centers of genetic diversity of quinoa (Apaza et al., 2015), and smallholders use five local commercial varieties, like Hualhuas, Junín white, Junín pink, Huancayo and Mantaro.

3.2 Choice experiment

3.2.1. Concept

We employ a discrete choice experiment to ex-ante analyze farmers' preferences and willingness to pay for an improved quinoa variety in the Junín region by presenting various hypothetical quinoa variety scenarios. Discrete choice experiments are a stated-preference method and are based on Lancaster's consumer model, in which respondents are presented with several choice sets, each containing a number of alternatives between which respondents are asked to choose. Each alternative is described by its attributes and each attribute takes one of several levels that describe ranges over which the attributes vary (Lancsar, Fiebig, & Hole, 2017). As a result, respondents derive utility from the different attributes of a good rather than the good as a whole (Lancaster, 1966). It is assumed that respondents choose the alternative which maximizes utility among the available options (McFadden, 1973). Experimental choice modelling allows to assess ex-ante preferences, when the good or service is not available in the market.

Choice experiments have become a popular tool to evaluate the preferences of respondents in marketing research (Alfnes, Guttormsen, Steine, & Kolstad, 2006; Loureiro & Umberger, 2007), labor economics (Ochieng, Veetil, & Qaim, 2017; Van den Broeck, Van Hoyweghen, & Maertens, 2016), agricultural technology studies (Asrat et al., 2010; Kassie et

al., 2017; Lambrecht et al., 2015; Ward et al., 2014), and studies on the adoption of sustainability standards (Ibnu, Glasbergen, Offermans, & Arifin, 2015; Meemken, Veetil, & Qaim, 2017). Several studies have used choice experiments to analyze the adoption of improved varieties among smallholder farmers in some staple food crop sectors, including a study on maize in Zimbabwe (Kassie et al., 2017), on rice and pearl millet in India (Smale et al., 2016; Ward et al., 2014), on beans in Burundi (Lambrecht et al., 2015), and on sorghum and teff in Ethiopia (Asrat et al., 2010).

3.2.2. Design

The design of a choice experiment entails different steps. The first step includes the identification of relevant attributes and attribute levels. Based on a review of the literature on quinoa seed, household survey data from 2014, semi-structured interviews with quinoa breeders at the National Institute of Agricultural Innovation (INIA), the Experimental Station of Santa Ana - Huancayo and agricultural experts at La Molina Agrarian University (UNALM), and four focus group discussions with quinoa farmers in the Junín region, we identify six attributes: 1) seed price, 2) grain size, 3) susceptibility to mildew, 4) maturation period, 5) yield and 6) saponin content (Table 1).

Table 1. Quinoa seed attributes and attribute levels used in the choice experiment

Attribute	Levels	Dummy coding
Seed price	4 PEN/kg	Continuous variable
	9 PEN/kg (SQ)	
	14 PEN/kg	
	24 PEN/kg	
	34 PEN/kg	
	40 PEN/kg	
Grain size	Small (SQ)	0
	Large	1
Susceptibility to Mildew	Susceptible (SQ)	MD1= 0 ; MD2=0
	Tolerance	MD1= 0 ; MD2=1
	Resistance	MD1= 1 ; MD2=0
Maturation period	4 months	Continuous variable
	5 months	
	6 months (SQ)	
	6 months (SQ)	
Yield	1 tons/ha	Continuous variable
	2 tons/ha (SQ)	
	3 tons/ha	
	4 tons/ha	
	5 tons/ha	
	6 tons/ha	
Saponin content	High level (bitter)	SC1= 1 ; SC2= 0
	Low content (semi-sweet) (SQ)	SC1= 0 ; SC2= 0
	Without saponin (sweet)	SC1= 0 ; SC2= 1

Note: SQ: Status quo

The first attribute is seed price, which is a monetary attribute and is expressed as the price of one kilogram of seed. The six levels (4, 9, 14, 24, 34 and 40 PEN/kg) were chosen based on

the average seed price in 2014 (about 9 PEN/kg), and the maximum and minimum price that focus group respondents found acceptable. We expect farmers to prefer a lower seed price.

The second attribute, grain size, describes the diameter of the grain expressed by two levels: 1) small grain with less than 1.8 mm, and 2) large grain with a diameter major of 1.8 mm. The current varieties that farmers use have a small grain size, which is more difficult to transform into pearled seeds or flakes by the food industry. We expect that farmers prefer a large grain size in order to receive a better price and sell to the food industry.

The third attribute, susceptibility to mildew, relates to the risk of yield loss when quinoa is affected by the mildew disease. We include three attribute levels: 1) resistance, with a risk of yield loss between 0% and 5%, 2) tolerance, with a risk of yield between 6% and 10% and 3) susceptible to mildew, with a risk of yield loss by more of 10%. We expect the farmer to dislike a susceptible quinoa seed.

The fourth attribute, maturation period, is expressed as the number of months between sowing and harvesting. Three levels (4, 5 and 6 months) have been chosen based on the minimum and maximum time to mature. If the maturation period is shorter, frost and drought are less likely to affect quinoa production and the opportunity costs of land (and labor) is reduced.

The fifth attribute, yield, is expressed as the expected output (in tons) of quinoa for one hectare. The yield levels (1, 2, 3, 4, 5 and 6 ton/ha) are based on the maximum level that experts deem to be feasible. The yield attribute is independent from the dosage of fertilizers and agricultural management. We expect that farmers have a strong preference for higher yields.

The sixth attribute, saponin content, is related to the level of saponin in the quinoa grain. We include three attribute levels: 1) bitter with a high level of saponin, 2) semi-sweet with low content of saponin, and 3) sweet without saponin. Saponin is found in the pericarp (outer shell) and gives the grain a bitter taste. It must be removed before the quinoa is edible, which increases the processing cost. We expect that farmers dislike seeds with higher saponin levels.

The second step of the design entails the construction of the choice cards. Each choice card includes two unlabeled alternatives that represent a hypothetical improved quinoa variety, described by the six attributes with varying levels. In addition, we include a status quo or opt-out option, indicating that the respondent is not willing to adopt an improved quinoa variety. If a farmer chooses this option, (s)he uses a white quinoa variety called Hualhuas that was cultivated by 78% of farmers in Junín region during the 2014 season. It is a long-maturation (6 months) variety yielding approximately 2 tons/ha under normal conditions. It is a small grain

with a low content of saponin (semi-sweet) and susceptible to mildew. The average seed price for this variety was about 9 PEN/kg in 2014.

Each farmer evaluated 12 choice cards. The total number of possible combinations of the two alternatives, the six attributes and its associated levels is much higher than 12, so we use a D-efficient design to reduce the number of choice cards to 12. This design aims to minimize the correlation between the attribute levels in the choice experiment and to minimize the standard errors of the parameter estimates. In a first phase, we use a small positive or negative number (0.001 and -0.001) to indicate whether we expect a positive or negative effect for a specific attribute. We expect that yield, grain size, mildew, and saponin content will positively influence the probability that an improved quinoa variety is chosen, while we expect a negative effect for maturity period and seed price. This preliminary design is used in a pilot test among 32 farmers who live in the Junín region. In a second phase, we use the parameter estimates from the pilot test in order to improve the efficiency of the design. The D-error of the preliminary design is 0.0933, while this is 0.0896 in the final design. We use Ngene software to generate the design. An example of a choice card is given in Figure A1.

3.2.3. Data collection

Data were collected in two phases. First, we collected baseline survey data between February and March 2015. We employed a three-stage sampling design with purposive selection in the first stage and stratified random selection in the second and third stage. In the first stage, based on statistics collected from Regional Direction of Agriculture of Junín (DRAJ, 2016), we identified 61 districts where quinoa was produced in 2014 and selected 25 that were more specialized in quinoa cultivation (with the median of the location ratios³ in the last three years more than one). In the second stage, based on reports and personal communication with employees from the agricultural agencies of Concepción, Chupaca and Jauja, we identified the largest villages in terms of population density (154 in total), from which we randomly sampled 47 villages. In the third stage, we randomly selected 518 farm-households, with the number of households in each village proportional to the population of quinoa producers. We used a quantitative structured questionnaire which provides household-level data on socio-demographic characteristics; land ownership; quinoa production and marketing; other crop

³ The location ratios ($Q_{ij} = (V_{ij} / \sum_i V_{ij}) / (\sum_j V_{ij} / \sum_i \sum_j V_{ij})$) represent the relationship between the participation of sector "i" in region "j" and the participation of the same sector in the national total and therefore, it is used as a measure of "relative or interregional specialization". The relative specialization of a region in an activity (sector) would be associated with $Q_{ij} > 1$; and it is measured with the harvested area between quinoa and total area cultivated (Lira, L., Quiroga, B., 2009).

production and marketing; livestock and animal production; off-farm activities and other income; and food security and quinoa consumption.

Second, we implemented a choice experiment in November and December 2016 using the same sample of farmers from the baseline survey. From the 518 farmers in the baseline sample, 458 farmers could be re-surveyed. This amounts to an attrition of 11.6%, which is due to local migration and refusal to participate in the survey. To ensure that the respondents understand the choice tasks, enumerators carefully explained the purpose of the choice experiment. They stressed that farmers should select the variety that they are most interested in, but that they should also be willing to pay the price indicated on the choice card. Each attribute and its levels were described into detail, and were visually represented on the choice cards. To test whether the choice experiment was understood by the farmers, we first presented a choice card in which there was one clearly dominant alternative before we proceeded with the actual choice experiment. In order to mitigate bias from respondents' fatigue or a learning effect, we presented the choice cards in a random order to each respondent. The choice experiment lasted about twenty minutes, and was accompanied by a small survey on current quinoa seed use and production/marketing practices.

3.2.4. Econometric analysis

To estimate farmers' preferences for different traits of improved quinoa varieties, we use a generalized multinomial logit (G-MNL) model (Fiebig, Keane, Louviere, & Wasi, 2010). The model incorporates both preference and scale heterogeneity, as it nests both the mixed logit (MXL) model and the scale heterogeneity multinomial logit (S-MNL) model. Preference heterogeneity relates to the fact that preferences for specific attributes are individual-specific and vary across respondents, while scale heterogeneity relates to the fact that choice consistency varies across respondents, implying an individual-specific error variance. Although preference heterogeneity has been widely acknowledged in the literature, there are few empirical studies that take scale heterogeneity into account (Kassie et al., 2017). The utility to farmer i ($i=1, \dots, 458$) from choosing alternative j ($j=1, 2, 3$) in choice card t ($t=1, \dots, 12$) in the generalized multinomial logit model (G-MNL) is given by:

$$U_{ijt} = \alpha_i ASC + \beta_i x_{ijt} + \varepsilon_{ijt}, \quad (1)$$

$$\beta_i = \sigma_i \beta + \gamma \eta_i + (1 - \gamma) \sigma_i \eta_i, \quad (2)$$

Where x_{ijt} is the vector of attributes; β_i is the vector of individual-specific parameters; and ε_{ijt} is the idiosyncratic error, which is assumed to be independently and identically distributed (i.i.d.) (eq.1). The specification of β_i (eq. 2) depends on β , which is a vector of mean

attribute utility weights; σ_i a person-specific scale heterogeneity of the idiosyncratic error which is log-normal with mean 1 and standard deviation τ , or $\text{LN}(1, \tau^2)$; η_i , which is a vector of individual specific deviations from the mean distributed $\text{MVN}(0, \Sigma)$; and γ , which is a scalar parameter that governs the variance of residual taste heterogeneity. Two model forms are produced to account for scale heterogeneity by restricting the scalar parameter γ . If γ is set to zero, the model G-MNL-I implies $\beta_i = \sigma_i \beta + \eta_i$, whereas if γ is set to 1, the model G-MNL-II implies $\beta_i = \sigma_i(\beta + \eta_i)$. The difference is that in the G-MNL-I model, the standard deviation of η_i is independent of the scaling of β , whereas in G-MNL-II, it is proportional to σ_i (Fiebig et al., 2010). However, if $\gamma < 0$ or $\gamma > 1$ we still have an interpretation (Keane & Wasi, 2013). Our model includes an alternative specific constant (ASC) to account for the alternative in which the farmer does not choose an improved variety but a traditional variety instead (meaning a positive α coefficient). The ASC is a dummy variable, coded zero for the improved varieties and one for the traditional variety. The ASC reflects a farmer's general attitude to the adoption of improved (quinoa) varieties, capturing factors not included by attributes in the choice experiment. We assume that farmers have a homogeneous preference for lower quinoa seed prices and we specify the price attribute to have a fixed coefficient. All other attributes and the ASC are specified as random and normally distributed. To account for preference heterogeneity, we estimate the G-MNL model for the full sample of farmers and for subsamples of food secure and food insecure farmers separately⁴ - with food security calculated as explained below. Assuming a linear utility function, the mean marginal willingness to pay (WTP) for a certain attribute x is given by the negative ratio of the coefficient for attribute x and cost (price of quinoa seed). This measure allows us to compare whether food secure farmers and food insecure farmers have different preferences for improved quinoa varieties.

3.3 Measurement of food security

We measure food security using the Food Insecurity and Hunger Module (FIHM) developed by the United States Department of Agriculture (USDA). Food security is defined as 'Access by all people at all times to enough food for an active, healthy life. It includes (1) the availability of nutritionally adequate and safe foods, and (2) the ability to acquire acceptable foods in socially acceptable ways' (Bickel, Nord, Price, Hamilton, & Cook, 2000). The FIHM is a questionnaire consisting of 18 multiple-choice questions, which measure the food security

⁴ We use this approach to identify preference heterogeneity instead of the more common approaches based on interaction terms in mixed logit models or latent class models as these approaches did not result in additional relevant insights.

situation at the household level during the last twelve months prior to the survey (Table A2). When a household experiences food insecurity, the frequency is assessed as well (ranging from ‘often’ to ‘some months but not every month’), which are coded as a positive response. Based on the sum of these responses, a score on the “Household Food Security Scale” is determined. This is a single numerical value ranging from 0 to 10, which measures the degree of severity of food insecurity/hunger experienced by a household. This measure allows us to classify households according to their food security status into four categories, each one representing a meaningful range of severity on the underlying scale: 1) food secure (from 0 to 2.2), 2) food insecure without hunger (from 2.4 to 4.4), 3) food insecure with moderate hunger (from 4.7 to 6.4), and 4) food insecure with severe hunger (from 6.6 to 10). We recode the household food security status as a binary variable ‘Food Insecure’, equal to one if a household is food insecure (irrespective of hunger status) and equal to zero if a household is food secure.

We acknowledge that other measures of food security, such as food consumption, energy and micronutrient intake, and anthropometric measures, have been shown to more correctly estimate food and nutrition insecurity than self-reported measures, like the FIHM (Lele et al., 2016). However, the FIHM has been demonstrated to be a useful, rapidly implementable, low-cost tool (MIDIS, 2012), which captures both physical and psychological elements of food insecurity.

4. Results

4.1. Descriptive statistics

Seventy percent of the sampled farm-households are classified as food secure and 30% as food insecure. The latter include households classified as ‘food insecure without hunger’ (21%), as ‘food insecure with moderate hunger’ (6%) and ‘food insecure with severe hunger’ (4%). Table 2 presents household and farm characteristics for the full sample, and across food secure and food insecure farm-households. Most (87%) of the sampled households are male headed and the average age of the household head is 50 years. While for food insecure households only 14% has a household head with completed tertiary education, this is 25% for food secure households. An average household consists of 3.7 members with food-insecure households having larger households. A small minority (19%) of farmers are member of a farmer organization, with food secure farmers being more likely to be member. In terms of household welfare, we find that the income of food secure farmers is more than triple the income of food insecure farmers, and that the propensity to be poor is 15% for food insecure farmers while this is only 3% for food secure farmers. These statistics confirm that households’ food security

status is highly correlated with welfare, which corroborates the validity of the food security measure.

Table 2. Household and farm characteristics by food security status

	Full sample		Food insecure households		Food secure households		
	Mean	SD	Mean	SD	Mean	SD	
Household head characteristics							
Age (year)	50.42	13.25	49.71	12.79	50.72	13.45	
Education level							
Secondary education	0.49		0.47		0.50		
Technical / university education	0.22		0.14		0.25		***
Female head (dummy)	0.13		0.11		0.14		
Household characteristics							
Household size total	3.74	1.52	4.03	1.50	3.62	1.51	***
Household size children	1.09	1.20	1.33	1.27	0.98	1.16	***
Organization member (dummy)	0.19		0.12		0.22		**
Net income 2014	20,435	45,242	8,414	14,068	25,619	52,515	***
Net income per capita	6,442	14,236	2,422	4,385	8,176	16,494	***
Net income per adult equivalent	10,084	21,672	4,042	7,048	12,690	25,079	***
Poor household MPI ⁵ (dummy)	0.07		0.15		0.03		***
Off-farm employment (dummy)	0.11		0.05	0.00	0.13	0.00	***
Farm characteristics							
Farm size (ha)	4.65	6.04	2.56	2.57	5.56	6.84	***
Number of crops	3.60	1.51	3.33	1.42	3.73	1.53	**
Number of tropical livestock units	3.42	5.24	3.04	5.66	3.58	5.05	
Quinoa area (ha)	1.85	2.88	1.02	1.19	2.20	3.30	***
Specialization ^a	0.43	0.24	0.44	0.24	0.42	0.24	
Quinoa price (PEN/kg)	6.86	2.20	6.49	2.45	7.02	2.07	***
Quinoa yield (kg/ha)	2,053	988	2,126	1,016	2,022	975	
Quinoa production (kg)	4,185	7,884	2,149	3,178	5,064	9,063	***
Self-consumption (%)	0.07	0.13	0.11	0.16	0.06	0.10	***
Seed-saving (%)	0.03	0.05	0.04	0.06	0.03	0.05	
Storage (%)	0.13	0.27	0.11	0.24	0.14	0.28	**
Sell (%)	0.76	0.31	0.75	0.30	0.76	0.31	
Experience growing quinoa (year)	10.51	11.09	8.29	8.79	11.43	11.80	***
Source of quinoa seeds							
from its last harvest (dummy)	0.47		0.43		0.49		
from other farmers (dummy)	0.46		0.51		0.44		
buying from INIA ^b (dummy)	0.05		0.06		0.05		
Use of Insecticide (dummy)	0.69		0.76		0.66		**
Use of Fungicide (dummy)	0.81		0.79		0.82		
Mildew problem (dummy)	0.82		0.78		0.84		
Extension on quinoa (dummy)	0.26		0.18		0.29		***

Note: Wilcoxon rank-sum test are used to test differences between food secure and food insecure households. Significant differences are indicated with * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. ^a Specialisation is measured as the

⁵ The Multidimensional Poverty Index (MPI) identifies multiple deprivations at the household level in health, education and standard of living and it was developed by the United Nations in 2010. Each household is assigned a deprivation score according to his deprivations in each of the 10 component indicators. The maximum deprivation score is 100 percent, with each dimension equally weighted. The health dimension has two indicators, nutrition and mortality, in which indicator is worth 16.7 percent. The education dimension has two indicators, years of schooling and child attendance to school, in with each indicator is worth 16.7 percent. The standard of living dimension has six indicators: cooking fuel, sanitation, water, electricity, floor and assets, in which indicator is worth 5.6 percent. Each household is classified as poor if the deprivation score is 33.3 percent or higher.

ratio of the surface devoted to quinoa area and its total surface cultivated, both measured in hectares. ^b INIA: National Institute of Agricultural Innovation.

Farmers in the research area cultivate on average 4.7 ha in total and allocate 1.9 ha to quinoa production. Food secure farmers produce more quinoa than food insecure farmers, but this is due to a larger farm size and not to higher yield levels, which is 2 tons/ha on average. They are also more likely to have received agricultural training and have been cultivating quinoa for a longer period (about three years more) than food insecure farmers. There are no differences related to quinoa seed use. The majority of farmers either store and reuse the harvested quinoa grain as seeds in the next season or buy it from other farmers. The five most common varieties are Hualhuas, Blanca de Junín, Rosada de Junín, Huancayo, and INIA 433 varieties, with Hualhuas as the most common variety (78% of farmers cultivating it during harvest 2014).

4.2. GMNL model results

Table 3 presents the empirical results of the generalized multinomial logit model (GMNL). The scale parameter τ is 0.6 with a standard error of 0.07, implying substantial scale heterogeneity in the data. We account for this in the models with gamma fixed at zero (GMNL I) and gamma fixed at one (GMNL II) (Table A3). The results of these models are very similar to the results of the unrestricted GMNL model, so we can assume that our results are robust to scale heterogeneity. We proceed the analysis with the unrestricted GMNL model and estimate this for food secure and food insecure farm-households separately.

Regarding the full sample, we find that the ASC estimate is negative and significant. This suggests that farmers generally prefer improved varieties over traditional varieties. All attribute coefficients are significant with expected signs (except for sweet quinoa, which is not significant). This suggests that all of these attributes matter in farmers' choice for improved varieties, and that farmers prefer cheaper varieties that are tolerant/resistant to mildew, and that have a larger grain size, higher yield levels and a shorter maturation period. Farmers dislike a high level of saponin but are indifferent between low levels of saponin and no saponin at all. However, we also find significant standard deviations for all of these attributes (except for tolerance to mildew and sweet quinoa), which suggests that preferences differ across farmers. We indeed find differences across food secure and food insecure farmers. The preferences for quinoa variety traits of food insecure farmers differ from those of food secure farmers with respect to maturation period and grain size, for which these farmers are indifferent.

Table 3. Generalized multinomial logit model estimates

	Full sample			Food insecure farmers			Food secure farmers		
	Mean	SE		Mean	SE		Mean	SE	
<i>Parameters</i>									
Seed price (PEN/kg)	-0.019	0.002	***	-0.027	0.009	***	-0.016	0.003	***
ASC	-0.724	0.109	***	-0.740	0.223	***	-0.741	0.138	***
Yield (ton)	0.971	0.060	***	0.991	0.224	***	0.946	0.122	***
Maturation period (months)	-0.061	0.027	**	-0.076	0.058		-0.063	0.032	**
Large grain ^a	0.184	0.060	***	0.207	0.227		0.189	0.104	*
Resistance to Mildew ^b	1.528	0.108	***	1.565	0.325	***	1.514	0.178	***
Tolerance to Mildew ^b	0.903	0.078	***	1.017	0.196	***	0.814	0.103	***
Bitter ^c	-3.308	0.222	***	-3.493	0.909	***	-3.194	0.451	***
Sweet ^c	-0.027	0.048		-0.106	0.098		0.012	0.058	
<i>Standard Deviations</i>									
ASC	1.231	0.084	***	1.145	0.205	***	1.298	0.112	***
Yield	0.465	0.047	***	0.410	0.151	***	0.409	0.050	***
Maturation period	-0.127	0.049	**	0.276	0.120	**	-0.079	0.063	
Large grain	0.638	0.088	***	0.773	0.422	*	0.608	0.150	***
Resistance to Mildew	0.364	0.120	***	-0.359	0.816		-0.363	0.165	**
Tolerance to Mildew	-0.012	0.072		0.178	0.322		-0.094	0.083	
Bitter	2.095	0.196	***	2.163	0.640	***	1.952	0.290	***
Sweet	0.129	0.082		0.064	0.366		-0.173	0.142	
τ	0.651	0.067	***	0.730	0.284	**	0.653	0.155	***
γ	-0.483	0.203	**	-0.022	0.353		-0.393	0.246	
Observations	16488			4968			11520		
Chi squared	386.641			66.485			155.771		
p-value	0.000			0.000			0.000		
Log likelihood	-4425.066			-1317.405			-3103.917		
BIC	9034.629			2796.514			6385.519		
AIC	8888.132			2672.809			6245.834		

Note: ASC, alternative specific constant (refers to traditional quinoa variety). ^a Reference category is a small grain. ^b Reference category is susceptible to Mildew. ^c Reference category is semi-sweet quinoa. * p < 0.1, ** p < 0.05, *** p < 0.01.

4.3. Willingness to pay for quinoa traits

The estimated WTP values for each attribute level are given in Table 4 for the full sample as well as for food secure and food insecure households. These WTP estimates highlight the extent to which farmers value characteristics of an improved quinoa variety. We find an average WTP of 49 PEN per kg of quinoa seed for a one ton higher yield and an average WTP of 81 PEN per kg of quinoa seed for mildew resistant varieties. These WTP values are high in comparison with current quinoa seed prices. According to the National Institute of Agricultural Innovation (INIA), the average price of quinoa seed during the time of our survey was 25 PEN/kg for a traditional variety (Hualhuas) and 40 PEN/kg for an improved variety (INIA 433/AIQ/FAO) in the Junín region. Given the discrepancy between current seed prices and the estimated WTP values, we need to interpret the WTP estimates cautiously and we do so in relative terms rather than in absolute terms.

Our results suggest that farmers are willing to pay most for an improved quinoa variety that is resistant to mildew. This is justified as mildew disease is the most important challenge

for quinoa production in the Junín region. Farmers are even prepared to pay double for a mildew-resistant variety than for a mildew-tolerant variety. Compared to the other traits, farmers value a mildew-resistant variety 1.6 times higher than a one-ton increase in yield level, 8.1 times higher than a variety with large grain size, and even 22.5 times higher than a one-month reduction in maturation period. However, farmers are not willing to buy a seed with a high saponin content, because the bitter taste does not allow for auto-consumption and makes it difficult to sell to food processors, as it increases their cost to remove the saponin.

Food secure farmers have the same ranking of attribute importance as food insecure farmers, but food secure farmers are in general willing to pay more for improved quinoa variety traits than food insecure farmers; e.g. 59.46% more for mildew-resistant varieties, 33.16% more for mildew-tolerant varieties, 55.83% more for a one-ton increase in yield level.

Table 4. Willingness to pay (WTP) estimates (PEN per kilogram of quinoa)

Attribute level	Total		Food insecure farmers		Food secure farmers		
	Mean	SD	Mean	SD	Mean	SD	
Yield	49.39	18.49	35.52	9.60	55.36	18.19	***
Maturation period	-3.62	3.31	-3.01 ^{n.s.}	5.50	-3.89	1.57	
Large grain	10.06	20.78	8.00 ^{n.s.}	18.02	10.95	21.83	
Resistance to Mildew	81.34	17.81	57.47	5.06	91.64	9.52	***
Tolerance to Mildew	46.02	5.90	37.37	1.72	49.76	1.48	***
Bitter	-172.73	87.69	-126.18	59.25	-192.81	90.36	***
Sweet	-0.84	3.44	-3.98 ^{n.s.}	0.50	0.52 ^{n.s.}	3.27	
Observations	458		138		320		

Note: WTP estimates were derived from generalized multinomial logit model parameter estimates. Wilcoxon rank-sum test are used to test differences between households with food security or not. Significant differences are indicated with * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. ^{n.s.} Not significant.

5. Discussion

In general, we find that traditional quinoa producers in the central highlands of Peru have a strong preference for improved varieties over traditional varieties – and a relatively high willingness to pay for improved quinoa varieties. Farmers have a strong preference for varieties that are resistant or tolerant to mildew. In addition, farmers prefer varieties that have higher yield levels, a larger grain size, and a shorter maturation period but their willingness to pay for these variety traits is substantially lower. Farmers dislike a high level of saponin. We find that farmers have heterogeneous preferences and that this preference heterogeneity can partially be explained by the food security status of the farm-household. Food insecure farmers, accounting for 30% of the sample, are found to be indifferent for maturation period and grain size, and to have a lower willingness to pay for improved varieties than food secure farmers. The indifference to maturation period and grain size likely relates to the fact that food insecure

farmers produce less quinoa and have a larger degree of self-consumption of quinoa (Table 2). As quinoa is less important as a cash crop for food insecure farmers, they do not appreciate a larger grain size, a characteristic that is valued in the market, nor a shorter maturation period, which may result in early marketing and better market prices and a lower opportunity cost of land. The lower willingness to pay likely relates to lower income levels, a higher incidence of poverty and higher capital constraints among food insecure farmers.

Mildew resistance is the most important attribute for the adoption of improved quinoa variety among smallholder farmers in Junín region. This is in line with the reported high incidence of mildew in Junín region (Rojas et al., 2015), causing yield reductions of up to 99% in susceptible cultivars (Zurita-Silva et al., 2014) and large income shocks. Our result that mildew resistance/tolerance, and hence yield stability, is more important to quinoa farmers than higher yield levels is consistent with findings in other studies – for example with the finding that drought tolerance is a more important trait than yield level for improved maize varieties in Zimbabwe (Kassie et al., 2017), and that farmers in Ethiopia are willing to forego a higher yield for more stability in yield with improved sorghum and teff varieties (Asrat et al., 2010). Our result that quinoa farmers prefer a shorter maturation period but that this trait is less important than mildew resistance and yield levels, is in line with a study on climbing beans in Burundi (Lambrecht et al., 2015) in which maturation period was found to be a less important trait than yield levels. Yet, other studies indicate stronger farmer preferences for yield levels than for yield variability (Lambrecht et al., 2015; Ward et al., 2014) – which implies that farmers' preferences for improved staple food crop varieties is context specific. We find substantial preference heterogeneity among farmers, which is in line with other choice experiment studies on staple food crop varieties (Asrat et al., 2010; Kassie et al., 2017; Lambrecht et al., 2015; Ward et al., 2014) that show preference heterogeneity related to food security status, resource endowments and risk aversion. A drawback of our study is that we were not able to explore preference heterogeneity more in depth, as we found few significant interactions with farm and farmer characteristics in the G-MNL model and derived no additional insights from latent class models. A novelty in our approach is the inclusion of scale heterogeneity, leading to results that are robust to heterogeneity in choice consistency across respondents – this source of heterogeneity was not accounted for in previous choice experiment studies on staple food crop varieties.

In addition, our results are consistent with other studies that consider high yields and lower seed prices as the main drivers for the adoption of improved varieties (Lambrecht et al., 2015; Sánchez-Toledano et al., 2017; Ward et al., 2014). For example, farmers had a strong

preference for rice and bean seed varieties that resulted in higher yields in India and Burundi respectively (Lambrecht et al., 2015; Ward et al., 2014). However, few studies that consider the local environmental conditions find that improved varieties with higher yield have a slight preference over the environmental adaptability (Asrat et al., 2010), and drought tolerant traits (Kassie et al., 2017) for a staple food crop like sorghum, teff and maize in Ethiopia and Zimbabwe, respectively.

In Peru, recent breeding programs focus on generating varieties with early maturity and short plant height for the introduction of quinoa in large-scale, modern and mechanized agricultural systems (Gomez-Pando & Eguiluz-de la Barra, 2013) – such as quinoa production in two seasons in high-technology farms along the Peruvian coast. However, small-scale and food insecure farmers who grow quinoa in non-irrigated and lowly mechanized systems in the highlands, do not benefit from these investments and improved varieties. From a public policy perspective, our results imply that developing mildew-resistant and higher-yielding varieties with a medium to low saponin content is a priority if investments in quinoa technologies are to benefit small-scale and food insecure farmers in the Andean highlands of Peru.

6. Conclusions

In this paper, we study farmers' preferences and willingness to pay for improved quinoa varieties using a choice experiment and data from 458 smallholder farmers in the Peruvian Andes. We find that farmers have a strong preference for improved quinoa varieties over traditional varieties, with mildew-tolerance or -resistance as most important crop trait. Farmers are found to prefer varieties that are characterized by higher yield levels, a larger grain size, lower levels of saponin (causing a bitter taste), and a reduced maturation period. Yet, food insecure farmers are found to be indifferent to early maturation and a larger grain size, which likely relates to quinoa being less important as a cash crop for these farmers – and have a lower willingness to pay for improved quinoa varieties. Our results imply that improved quinoa varieties entail the highest potential for a rapid diffusion among smallholder Andean farmers if ongoing investments focus on the development of mildew-resistant and higher-yielding varieties with a medium to low saponin content.

This paper contributes to the literature on ex ante adoption of improved varieties for staple food crops with new evidence on smallholders' preferences for quinoa varieties. Our study underscores the importance of studying farmers' preferences ex ante to better inform agricultural research and development initiatives – which holds especially in the case of quinoa,

given the sharp increase in attention for and investment in this crop in recent years – and corroborates earlier findings on the heterogeneity of preferences among farmers.

Appendix

Table A1. Quinoa harvested area, production, yield and consumption in quinoa-producing farmers in Peru.

	2010		2011		2012		Average
	Mean	SD	Mean	SD	Mean	SD	
N° farmers	424		354		573		
Area harvested (has)	0.21	0.53	0.16	0.38	0.19	0.44	0.18
Yield (kg/ha)	1,770	2,250	1,156	1,066	1,026	1,736	1,317
Production (kg)	228.91	763.98	163.56	489.56	234.51	1109.90	208.99
% of sell (Sale/Production)	0.18	0.30	0.17	0.30	0.15	0.29	0.16
% of Self-consumption (Self-consumption/Production)	0.56	0.31	0.60	0.33	0.60	0.34	0.59
Price (soles/kg)	3.71	1.02	3.74	0.84	4.12	1.10	3.86
Consumption per capita	19.98	41.93	15.02	18.72	15.70	22.33	16.90
Consumption per adult equivalent	31.63	71.48	20.65	24.94	21.40	27.74	24.56

Source: Author's calculation based on micro data from National Survey of Strategic Programs (ENAPRES), over 2010-2012

Table A2. Food Insecurity and Hunger Module (FIHM)


Question description	Yes	No	Percentage (N=458)			
			Often	Sometimes	Never	Refused
Q2 Worried whether food would run out	--	--	3.71	47.82	46.51	1.97
Q3 Food that we bought just did not last	--	--	1.75	34.72	61.79	1.75
Q4 Could not afford to eat balanced meals	--	--	1.09	22.93	75.11	0.87
Q5 Relied on only a few kinds of low-cost food to feed children	--	--	0.66	13.54	84.93	0.87
Q6 Could not feed the children a balanced meal	--	--	0.44	10.04	88.65	0.87
Q7 Children were not eating enough	--	--	0.22	6.33	92.58	0.87
Q8* Adult cut the size of meals or skipped them	14.41	85.59	0.44	12.45	85.59	1.53
Q9 Eat less than felt should	16.38	83.62	--	--	--	--
Q10 Hungry but did not eat	12.88	87.12	--	--	--	--
Q11 Lose weight	12.45	87.55	--	--	--	--
Q12* Adult did not eat for a whole day	6.77	93.23	0.22	5.02	93.23	1.53
Q13 Cut the size of children's meals	5.02	94.98	--	--	--	--
Q14* Children ever skip meals	3.28	96.72	0.00	2.40	96.72	0.87
Q15 Children ever hungry	2.40	97.60	--	--	--	--
Q16 Children did not eat for a whole day	2.18	97.82	--	--	--	--

Note: For questions 8, 12 and 14, “Often” is “Almost every month” and “Sometimes” is “Some months but not every month”.

Table A3. Generalized multinomial logit model

	Full GMNL			GMNL- II			GMNL- I		
	Mean	SE		Mean	SE		Mean	SE	
<i>Parameters</i>									
ASC	-0.69	0.10	***	-0.71	0.11	***	-0.71	0.10	***
Seed price	-0.02	0.00	***	-0.02	0.00	***	-0.02	0.00	***
Yield	0.88	0.06	***	0.84	0.05	***	0.76	0.04	***
Maturation period	-0.07	0.03	***	-0.06	0.03	**	-0.05	0.02	*
Large grain	0.16	0.06	***	0.15	0.06	***	0.12	0.05	**
Resistance	1.40	0.10	***	1.38	0.10	***	1.23	0.07	***
Tolerance	0.83	0.07	***	0.80	0.07	***	0.72	0.06	***
Bitter	-3.08	0.22	***	-2.96	0.17	***	-2.64	0.14	***
Sweet	-0.03	0.04		-0.03	0.05		-0.01	0.04	
<i>Standard Deviations</i>									
ASC	1.22	0.12	***	1.23	0.10	***	1.19	0.10	***
Yield	0.39	0.04	***	0.39	0.04	***	0.36	0.03	***
Maturation period	-0.14	0.05	***	-0.15	0.07	**	-0.19	0.04	***
Large grain	0.55	0.11	***	0.41	0.10	***	0.52	0.08	***
Resistance	-0.21	0.16		-0.29	0.11	***	0.19	0.28	
Tolerance	0.06	0.08		0.11	0.10		0.19	0.12	
Bitter	1.82	0.19	***	1.75	0.15	***	1.49	0.11	***
Sweet	0.07	0.11		0.01	0.23		-0.08	0.09	
Tau	-0.60	0.09	***	0.57	0.08	***	0.35	0.08	***
Gamma	-0.24	0.18							
Observations	16488			16488			16488		
Chi squared	343.11			465.41			817.56		
p-value	0.00			0.00			0.00		
Log likelihood	-4452.19			-4444.82			-4464.48		
AIC	8942.378			8925.647			8964.966		
BIC	9088.875			9064.434			9103.753		

Note: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$



Usted prefiere la alternativa **A**, **B** o su **Situación Actual (C)**

Figure A1. Example of a choice card N° 1 as shown to quinoa producers during the interview.

Note: Authors' elaboration.

References

- Alfnes, F., Guttormsen, A. G., Steine, G., & Kolstad, K. (2006). Consumers' Willingness to Pay for the Color of Salmon: A Choice Experiment with Real Economic Incentives. *Amer. J. Agr. Econ.*, 88(4), 1050–1061.
- Apaza, V., Cáceres, G., Estrada, R., & Pinedo, R. (2015). *Catalogue of commercial varieties of quinoa in Perú*.
- Asrat, S., Yesuf, M., Carlsson, F., & Wale, E. (2010). Farmers' preferences for crop variety traits: Lessons for on-farm conservation and technology adoption. *Ecological Economics*, 69(12), 2394–2401.
- Bazile, D., Bertero, D., & Nieto, C. (2015). *State of the Art Report on Quinoa around the World in 2013* (Rome).
- Bazile, D., Jacobsen, S.-E., & Verniau, A. (2016). The Global Expansion of Quinoa: Trends and Limits. *Frontiers in Plant Science*, 7, 1–6.
- Bazile, D., Pulvento, C., Verniau, A., Al-Nusairi, M. S., Ba, D., Breidy, J., ... Padulosi, S. (2016). Worldwide Evaluations of Quinoa: Preliminary Results from Post International Year of Quinoa FAO Projects in Nine Countries. *Frontiers in Plant Science*, 7.
- Bedoya-Perales, N. S., Pumi, G., Mujica, A., Talamini, E., & Padula, A. D. (2018). Quinoa expansion in Peru and its implications for land use management. *Sustainability (Switzerland)*, 10(2), 1–13.
- Bickel, G., Nord, M., Price, C., Hamilton, W., & Cook, J. (2000). *Guide to Measuring Household Food Security Revised 2000*. Agriculture.
- Bojanic, A. (2011). *La quinua : Cultivo milenario para contribuir a la seguridad alimentaria mundial*. Proinpa.
- Danielsen, S., & Munk, L. (2004). Evaluation of disease assessment methods in quinoa for their ability to predict yield loss caused by downy mildew. *Crop Protection*, 23(3), 219–228.
- DRAJ. Regional Direction of Agriculture of Junin. (n.d.). Time Series of Agricultural Production. Retrieved June 3, 2018, from http://www.agrojunin.gob.pe/?page_id=356
- Fiebig, D. G., Keane, M. P., Louviere, J., & Wasi, N. (2010). The Generalized Multinomial Logit Model: Accounting for Scale and Coefficient Heterogeneity. *Marketing Science*, 29(3), 393–421.
- Gomez-Pando, L. R., & Eguiluz-de la Barra, A. (2013). Developing Genetic Variability of Quinoa (*Chenopodium quinoa* Willd.) with Gamma Radiation for Use in Breeding Programs. *American Journal of Plant Sciences*, 04(02), 349–355.
- Hellin, J., & Higman, S. (2005). Crop diversity and livelihood security in the andes.

Development in Practice, 15(2), 165–174.

- Ibnu, M., Glasbergen, P., Offermans, A., & Arifin, B. (2015). Farmer Preferences for Coffee Certification: A Conjoint Analysis of the Indonesian Smallholders. *Journal of Agricultural Science*, 7(6), 20–35.
- INEI. National Institute of Statistics and Informatics. (n.d.). National Survey of Strategic Programs (ENAPRES). Retrieved January 3, 2018, from <http://iinei.inei.gob.pe/microdatos/>
- Jacobsen, S.-E. (2003). The Worldwide Potential for Quinoa (*Chenopodium quinoa* Willd.). *Food Reviews International*, 19(1–2), 167–177.
- Jacobsen, S.-E., Mujica, A., & Jensen, C. R. (2003). The Resistance of Quinoa (*Chenopodium quinoa* Willd.) to Adverse Abiotic Factors. *Food Reviews International*, 19(1–2), 99–109.
- Jacobsen, S. E., Mujica, A., & Ortiz, R. (2003). The Global Potential for Quinoa and Other Andean Crops. *Food Reviews International*, 19(1–2), 139–148.
- Kassie, G. T., Abdulai, A., Greene, W. H., Shiferaw, B., Abate, T., Tarekegne, A., & Sutcliffe, C. (2017). Modeling Preference and Willingness to Pay for Drought Tolerance (DT) in Maize in Rural Zimbabwe. *World Development*, 94, 465–477.
- Keane, M., & Wasi, N. (2013). Comparing alternative models of heterogeneity in consumer choice behavior. *Journal of Applied Econometrics*, 28, 1018–1045.
- Lambrecht, I., Vranken, L., Merckx, R., Vanlauwe, B., & Maertens, M. (2015). Ex ante appraisal of agricultural research and extension: A choice experiment on climbing beans in Burundi. *Outlook on Agriculture*, 44(1), 61–67.
- Lancaster, K. (1966). A New Approach to Consumer Theory. *Chicago Journals*, 74(2), 132–157.
- Lancsar, E., Fiebig, D. G., & Hole, A. R. (2017). Discrete Choice Experiments: A Guide to Model Specification, Estimation and Software. *PharmacoEconomics*, 35(7), 697–716.
- Lele, U., Masters, W. A., Kinabo, J., Meenakshi, J. V, Ramaswami, B., Tagwireyi, J., & Goswami, S. (2016). *Measuring Food and Nutrition Security : An Independent Technical Assessment and User's Guide for Existing Indicators*. Rome.
- Lester, G. E. (2006). Organic versus conventionally grown produce: Quality differences, and guidelines for comparison studies. *HortScience*, 41(2), 296–300.
- Loureiro, M. L., & Umberger, W. J. (2007). A choice experiment model for beef: What US consumer responses tell us about relative preferences for food safety, country-of-origin labeling and traceability. *Food Policy*, 32(4), 496–514.
- McFadden, D. (1973). Conditional logit analysis of qualitative choice behavior. *Frontiers in*

Econometrics.

- Meemken, E. M., Veettil, P. C., & Qaim, M. (2017). Toward Improving the Design of Sustainability Standards—A Gendered Analysis of Farmers' Preferences. *World Development*, 99, 285–298.
- MIDIS. (2012). *Mapa de vulnerabilidad a la inseguridad alimentaria 2012*.
- MINAGRI. Ministry of agriculture and Irrigation. (n.d.). Time Series of Agricultural Production. Retrieved March 10, 2018, from http://frenteweb.minagri.gob.pe/sisca/?mod=consulta_cult
- Mujica, A. (1992). Granos y leguminosas andinas. In J. . Hernández Bermejo & J. León (Eds.), *Cultivos Marginados: otra perspectiva de 1492* (pp. 129–146). FAO.
- Ochieng, D. O., Veettil, P. C., & Qaim, M. (2017). Farmers' preferences for supermarket contracts in Kenya. *Food Policy*, 68, 100–111.
- Ofstehage, A. (2012). The construction of an alternative quinoa economy: Balancing solidarity, household needs, and profit in San Agustín, Bolivia. *Agriculture and Human Values*, 29(4), 441–454.
- Rojas, W., Pinto, M., Alanoca, C., Gómez Pando, L., Leon-Lobos, P., Alercia, A., ... Bazile, D. (2015). Quinoa genetic resources and ex situ conservation. In *State of the Art Report on Quinoa Around the World in 2013* (pp. 56–82).
- Ruiz, K. B., Biondi, S., Oses, R., Acuña-Rodríguez, I. S., Antognoni, F., Martínez-Mosqueira, E. A., ... Molina-Montenegro, M. A. (2014). Quinoa biodiversity and sustainability for food security under climate change. A review. *Agronomy for Sustainable Development*, 34(2), 349–359.
- Sánchez-Toledano, B. I., Kallas, Z., & Gil-Roig, J. M. (2017). Farmer preference for improved corn seeds in Chiapas, Mexico: A choice experiment approach. *Spanish Journal of Agricultural Research*, 15(3).
- Sibiya, J., Tongoona, P., Derera, J., & Makandaa, I. (2013). Farmers' desired traits and selection criteria for maize varieties and their implications for maize breeding: A case study from Kwazulu-Natal Province, South Africa. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*, 114(1), 39–49.
- Smale, M., Diressie, M. T., & Birol, E. (2016). Understanding the potential for adoption of high-iron varieties of pearl millet in Maharashtra, India: what explains their popularity? *Food Security*, 8(2), 331–344.
- Stevens, A. W. (2017). Quinoa quandary: Cultural tastes and nutrition in Peru. *Food Policy*, 71(August), 132–142.

- Tapia, M. E., & Fries, A. M. (2007). *GUÍA DE CAMPO DE LOS CULTIVOS ANDINOS*.
- Van den Broeck, G., Van Hoyweghen, K., & Maertens, M. (2016). Employment Conditions in the Senegalese Horticultural Export Industry: A Worker Perspective. *Development Policy Review*, 34(2), 301–319.
- Ward, P. S., Ortega, D. L., Spielman, D. J., & Singh, V. (2014). Heterogeneous demand for drought-tolerant rice: Evidence from Bihar, India. *World Development*, 64, 125–139.
- Zurita-Silva, A., Fuentes, F., Zamora, P., Jacobsen, S. E., & Schwember, A. R. (2014). Breeding quinoa (*Chenopodium quinoa* Willd.): Potential and perspectives. *Molecular Breeding*, 34(1), 13–30.